

THE UNCERTAINTY PRINCIPLE FOR OPERATORS DETERMINED BY LIE GROUPS

JENS GERLACH CHRISTENSEN

ABSTRACT. For unbounded operators A, B and C in general, the commutation relation $[A, B] = C$ does not lead to the uncertainty relation $\|Au\|\|Bu\| \geq \frac{1}{2}|\langle Cu, u \rangle|$. If A, B and C are part of the generators of a unitary representation of a Lie group then the uncertainty principle above holds.

If A and B are self-adjoint or skew-adjoint operators then

$$(1) \quad \|Au\|\|Bu\| \geq \frac{1}{2}|\langle [A, B]u, u \rangle| \quad \text{for all } u \in D([A, B])$$

The operator $[A, B]$ need not be closed, and if $C = \overline{[A, B]}$, the extended inequality

$$(2) \quad \|Ax\|\|Bx\| \geq \frac{1}{2}|\langle Cx, x \rangle| \quad \text{for all } x \in D(A) \cap D(B) \cap D(C).$$

need not be valid. K. Kraus [4] showed that (2) holds when A, B and C are infinitesimal generators of a unitary representation of a Lie group of dimension ≤ 3 . The result of K. Kraus was slightly extended by G.B. Folland and A. Sitaram in [1, Theorem 2.4], but the necessity of the dimension constraint was left as an open problem. In the present paper the dimension constraint is shown to be unnecessary, by means of some results by I.E. Segal in [2].

If (H, π) is a unitary representation of a Lie group G and R is a right-invariant vector field, then by Stone's theorem one can define the closed skew-adjoint (by Lemma 3.1.13 in [2]) operator $\pi(R)$ such that $\pi(\exp tR) = \exp(\pi(tR))$ for all $t \in \mathbb{R}$. Then $[\pi(X), \pi(Y)] \subseteq \pi([X, Y])$.

Theorem. *Let G be a Lie group with Lie algebra $\mathfrak{g}$, and let (H, π) be a unitary representation of G . Suppose that $X, Y \in \mathfrak{g}$. Then*

$$\|Ax\|\|Bx\| \geq \frac{1}{2}|\langle Cx, x \rangle| \quad \text{for all } x \in D(A) \cap D(B) \cap D(C)$$

holds with $A = \pi(X), B = \pi(Y)$ and $C = \pi([X, Y])$.

Lemma. *Let σ be the modular function on G . Given a right-invariant vector field R on G , let L be the corresponding left-invariant vector field, and let $\gamma = \frac{d}{dt}\sigma(\exp(tR))|_{t=0}$.*

- (a) *If $f \in C_c^\infty(G)$ and $f^*(g) = \overline{f(g^{-1})}\sigma(g)$, then $\pi(Rf^*)^* = \pi(Lf) + \gamma\pi(f)$.*
- (b) *There is a sequence $\{f_n\}$ in $C_c^\infty(G)$ such that $\pi(f_n)u \rightarrow u$ and $\pi(Rf_n)u + \pi(Lf_n)u + \gamma\pi(f_n)u \rightarrow 0$ for all $u \in H$.*

2000 *Mathematics Subject Classification.* Primary 22E45, 47B15, 47B25, 81Q10.

The author wishes to thank Henrik Schlichtkrull for his proofreading and kind supervision.

Proof. (a) is simply Lemma 3.1.7 in [2] using the fact that $f^{**} = f$. (b) is Lemma 3.1.8 and Lemma 3.1.9 in [2], and $\pi(f_n)u \rightarrow u$ has been proved in [3, p.56]. $\square$

To prove the theorem I will first show that for $R \in \mathfrak{g}$ it holds that $\pi(R)\pi(f_n)u \rightarrow \pi(R)u$ for $u \in D(\pi(R))$. Fix $R \in \mathfrak{g}$ and let L be the left-invariant vector field corresponding to R . Note that for all $f \in C_c^\infty(G)$ it holds that $\pi(f)\pi(R) \subseteq -\pi(Rf^*)^*$, since $\langle \pi(f)\pi(R)u, v \rangle = \langle u, -\pi(Rf^*)v \rangle$ for all $u \in D(\pi(R))$ and $v \in H$. Hence

$$\pi(f)\pi(R) \subseteq -\pi(Lf) - \gamma\pi(f)$$

by (a) of the lemma. But then

$$\pi(R)\pi(f_n)u - \pi(f_n)\pi(R)u = \pi(Rf_n)u + \pi(Lf_n)u + \gamma\pi(f_n)u$$

for any $u \in D(\pi(R))$. By (b) in the lemma this tends to 0 and since $\pi(f_n)\pi(R)u \rightarrow \pi(R)u$ it follows that $\pi(R)\pi(f_n)u \rightarrow \pi(R)u$.

Now let A, B and C be as in the theorem. For any $u \in D(A) \cap D(B) \cap D(C)$ it follows that $\pi(f_n)u \rightarrow u$, $A\pi(f_n)u \rightarrow Au$, $B\pi(f_n)u \rightarrow Bu$ and $C\pi(f_n)u \rightarrow Cu$. Since the Gårding vector $\pi(f_n)u$ is in $D([A, B])$ by [3, p.56] the inequality (1) gives (2) as desired.

REFERENCES

- [1] Gerald B. Folland and Alladi Sitaram, *The Uncertainty Principle: A Mathematical Survey*, The Journal of Fourier Analysis and Applications, Vol. 3, Number 3, 207–238, 1997
- [2] I. E. Segal, *A class of operator algebras which are determined by groups*, Duke Math. Journal, Vol. 18, 221–265, 1951
- [3] Anthony W. Knap, *Representation Theory of Semisimple Groups*, Princeton University Press, Princeton, New Jersey, 1986
- [4] K. Kraus, *A Further Remark on Uncertainty Relations*, Zeitschrift für Physik, 201, 134–141, 1967

MATEMATISK INSTITUT, KØBENHAVNS UNIVERSITET

Current address: Universitetsparken 5, 2100 København Ø, Denmark

E-mail address: vepjan@math.ku.dk

URL: <http://www.math.ku.dk/~vepjan>